

# **Terminal Restriction Fragmentation Length Polymorphism Analysis of 16S rRNA Genes for the Characterization of Bacterial Community Structure in the Rumen of Sheep Fed Different Diets**

Eiichi MIYAGAWA, Jyunki OKUYAMA, Yutaka HANDA, Kazuo SATO  
Masateru KOIWA and Eiryu OKAMOTO

酪農学園大学紀要 別刷 第31巻 第2号

*Reprinted from*

"Journal of Rakuno Gakuen University" Vol.31, No.2 (2007)

## Terminal Restriction Fragmentation Length Polymorphism Analysis of 16S rRNA Genes for the Characterization of Bacterial Community Structure in the Rumen of Sheep Fed Different Diets

Eiichi MIYAGAWA, Jyunki OKUYAMA, Yutaka HANDA, Kazuo SATO<sup>1</sup>,  
Masateru KOIWA<sup>2</sup> and Eiryu OKAMOTO  
(October 2006)

### Introduction

The dependence of rumen bacterial community structure on an animal's diet is a well-documented fact (Dehority and Orpin, 1977). Until recently, any changes in an animal's bacterial community have been monitored essentially using culture-dependent techniques. However, even when the largest number of colonies grew on a medium by these techniques, it usually only reached about one-tenth the number determined by the direct counting method (Suto, 1970; Minato et al., 1990). Furthermore, analysis of these microbial communities by culture methods requires a great deal of labor and time. Minato et al. (1990) proposed non-cultivation based techniques for investigating rumen bacterial communities, that is, method which employ counting, fractionation and chemotaxonomy. But there have also been limitations in clarifying microflora in rumen using these techniques. Recently, the molecular biological methods have brought new developments into the field, omitting the cultivation step. Molecular biological analysis techniques which have been used to analyze rumen microbial communities include: dot blot hybridization (Attwood et al., 1988; Stahl et al., 1988), sequence analysis of 16S rDNA libraries (Whitford et al., 1998; Tajima et al., 1999; Kocherginskaya et al., 2001), denaturing

gradient gel electrophoresis (DGGE) (Kocherginskaya et al., 2001), fluorescence *in situ* hybridization (FISH) (Yanagita et al., 2000), competitive PCR (Koike et al., 2003) and real-time PCR (Tajima et al., 2001).

Terminal restriction fragment length polymorphism (T-RFLP), which is also a molecular biological method, is a PCR-based method for rapidly comparing bacterial communities independent of culture or cloning (Kitts, 2001). The method was first applied to a mixture of different pure cultures of bacteria and was found to be useful for defining the number of operational taxonomic units (OTUs) present (Avaniss-Aghajani et al., 1994). Following this, T-RFLP analysis has been implemented in the characterization of microbial diversity in activated sludge (Hiraishi et al., 2000), soil (Blackwood et al., 2003; Dunber et al., 2000), human feces (Hayashi et al., 2002; Nagashima et al., 2003), the colon of pigs (Leser et al., 2000), marine sediment (Scala and Kerkhof, 2000), oral bacterial flora (Sakamoto et al., 2003) and so on. T-RFLP analysis provides size in base pairs about each of the terminal restriction fragments (T-RF) detected. T-RF sizes can be compared to a database of theoretical T-RFs derived from sequence information. Since T-RFLP patterns also provide quantitative information on the relative abundance of T-RFs, it is easier to process

---

酪農学園大学酪農学部酪農学科農業微生物学研究室

Agricultural Microbiology, Department of Dairy Science, Faculty of Dairy Science, Rakuno Gakuen University, 582 Ebetsu, Hokkaido, 069-8501, Japan

<sup>1</sup> 酪農学園大学酪農学部酪農学科酪農経営学研究室

Dairy Management, Department of Dairy Science, Rakuno Gakuen University, 582 Ebetsu, Hokkaido, 069-8501, Japan

<sup>2</sup> 酪農学園大学獣医学部獣医学科生産動物医療学教室

Animal Medical Care (Large Animal), Department of Veterinary Medicine, Rakuno Gakuen University, 582 Ebetsu, Hokkaido, 069-8501, Japan

data by appropriate numerical and statistical methods. Little has been reported on T-RFLP analysis in rumen bacterial ecosystems. The authors herein report the use of T-RFLP as a means to compare the bacterial community structure in the rumen of sheep fed different diets.

### Materials and Methods

**Animals and feeds.** Three Suffolk sheep (A, B, and C), weighing between 69-87 kg, and each fitted with a rumen fistula, were used. Before the experiments, animals were maintained on a common diet consisting of 1.0 kg of timothy hay and 0.1 kg of mixed feed ( $\alpha$ Merit 16; Chubu Shiryō Co., Ltd.), being fed once a day. The sheep were reared in individual pens (Sanshin Industrial Co.) in a sheep house at Rakuno Gakuen University. The feed was changed from the mixture of common feed and timothy hay to solely timothy hay (100%). After animals were maintained on the roughage for 10 days, the experiment was performed on the 11th day. Following this, the diet was changed from the timothy hay (100%) to a high concentrate diet (timothy hay and barley pellets; 40%:60%, w/w), increasing the amount of barley little by little to allow time (for 26 days) for adaptation. Timothy hay was cut in about 10 cm lengths by a cutter, and the barley pellets were mashed with a mixer. The high concentrate diet was prepared by mixing cut timothy and mashed barley pellets (40%:60%, w/w) with water at a ratio of 0.7-0.8/1 (wt/wt). After the sheep were fed this diet for 10 days, the experiment was performed on the 11th day. The amount of both the roughage and high concentrate diets supplied once daily were about 1.5 % per body weight. Mineral (Nippon Formula Feed Mfg Co.) and water were available *ad libitum*.

**Preparation of rumen bacterial fraction.** About 10 ml of the rumen contents were sampled using a stomach tube before the morning feed, and every two hours after feeding. The rumen contents were then squeezed through two layers of cheesecloth. The rumen bacterial fraction from the filtrate were harvested by centrifugation (17,400 $\times$ g, 10 min) at 4°C, washed with TE buffer (10mM Tris-HCl, 1 mM Na<sub>2</sub>EDTA, pH8.0) until

the color of the supernatant disappeared. The precipitate was kept at -20°C before use. Bovine rumen content obtained from a slaughterhouse located at Hayakita-cho in Hokkaido was used for the selection test of restriction enzymes for T-RFLP analysis.

**Rumen fluid components.** The pH of the rumen fluids was measured with a pH meterΦ260 (Beckman, U. S. A.) immediately after collection. Volatile fatty acids (VFAs) and non-VFAs were analyzed using a gas chromatograph (Shimadzu GC-8A) equipped with a flame ionization detector with N<sub>2</sub> as the carrier gas, by injecting 1 $\mu$ l of deproteinized sample with 24% meta-phosphoric acid in 5N-H<sub>2</sub>SO<sub>4</sub> (Ueki et al., 1978). The ammonium concentration of rumen fluids was measured by the method of Chaney and Marbach (1962), using NH<sub>4</sub>Cl as the standard.

**DNA extraction from the rumen bacterial fraction.** Total DNA was extracted from the rumen bacterial fraction according to the protocol previously described by Hiraishi et al. (2000) with minor modifications. The frozen sample (0.1-0.2g; wet weight) was thawed at room temperature and suspended in 1 ml of TE buffer, to which 10 $\mu$ l of lysozyme solution (1 mg/ml; Wako Pure Chemical Industries, Ltd.) was added. The suspension was subjected to 2 min disruption with a beads homogenizer (CSC model BC-20, Central Scientific Commerce, Inc.). The suspension was centrifuged at 2,400 $\times$ g for 5 min with a high speed refrigerated micro centrifuge (Tomy MX-150:4°C). The pellet was resuspended in TE buffer and disrupted two times in the same way. These three successive supernatants (about 3 ml) were combined. Ten  $\mu$ l of proteinase K (10mg/ml; Wako) was added to 1 ml of the supernatant in a 1.5 ml Eppendorf tube, and incubated at 55°C for 30 min. Fifty  $\mu$ l of 10 % sodium dodecyl sulfate solution was put into the tube and reacted at 60°C for 30 min. After treatment with hexadecyltrimethyl ammonium bromide, the supernatant was treated with phenol-chloroform-isoamyl alcohol (25:24:1, v/v) which was shaken for a few minutes, and centrifuged at 13,900 $\times$ g for 5 min. Twofold volumes of isopropyl alcohol and one tenth volumes of 3 M sodium acetate (pH

5.2) were added to the supernatant, and allowed to stand overnight at  $-20^{\circ}\text{C}$ . The solution was centrifuged at  $17,400\times g$  for 30 min ( $4^{\circ}\text{C}$ ), and the resulting supernatant was removed. One ml of 70% ethyl alcohol was added to the residue, and re-centrifuged at  $17,400\times g$  for 10 min. After air-drying the precipitate, the purified DNA was dissolved in  $50\text{ }\mu\text{l}$  of TE buffer by shaking at  $65^{\circ}\text{C}$  for a few minutes using a shaker.

**PCR amplification of 16S rDNA.** 16S rDNA fragments were amplified by polymerase chain reaction (PCR) with universal primers 27f (5'-AGAGTTTGATCCTGGCTCAG-3') and 1492r (5'-GGCTACCTTGTACGACTT-3') (Lane, 1991) which corresponds to position 8 to 27 and 1510 to 1492, respectively, in *Escherichia coli* 16S rRNA (Brosius et al., 1978). These primers were purchased from Amersham-Pharmacia Biotech KK (Tokyo). PCR was performed with 0.5 ng of the purified DNA as the template. Thirty nine  $\mu\text{l}$  of distilled water,  $200\mu\text{M}$  (each) of deoxynucleoside triphosphates,  $0.2\mu\text{M}$  of both primers, and 1.25 U of Ex *Taq* DNA polymerase (Takara Shuzo Co., Kyoto) were included in the  $50\text{ }\mu\text{l}$  reaction mixture. The PCR was conducted with a thermal cycler (ABI GeneAmp PCR System 9700, Applied Biosystems). PCR cycling consisted of an initial denaturation at  $93^{\circ}\text{C}$  for 3 min; followed by 30 cycles of denaturation at  $93^{\circ}\text{C}$  for 30 sec, annealing at  $53.5^{\circ}\text{C}$  for 30 sec, and extension at  $72^{\circ}\text{C}$  for 90 sec; and a final extension at  $72^{\circ}\text{C}$  for 7 min. The PCR products were purified according to the manufacturer's instructions using a QIAquick PCR purification kit (Qiagen Co., Tokyo). The forward primer labeled with 5'-Fam (carboxy-fluorescein-*N*-hydroxysuccinimide ester-dimethyl sulfoxide), designated Fam-27f and the reverse primer 907r (5'-CCGTCAATTCATTTGAGTTT-3' (Hiraishi et al., 2000) were used in the next nested PCR. The cycle profiles were denaturation at  $95^{\circ}\text{C}$  for 5min; followed by 30 cycles of denaturation at  $95^{\circ}\text{C}$  at 30 sec, annealing at  $55^{\circ}\text{C}$  for 30 sec, and extension at  $72^{\circ}\text{C}$  for 1 min; and final extension at  $72^{\circ}\text{C}$  for 7 min. The PCR products were purified using the PCR purification kit.

**T-RFLP analysis.** Three tandem tetrametric

restriction endonucleases, *HhaI*, *MspI*, and *AfaI*, were used for cutting the 16S rDNA fragments. The purified PCR products were cut with each of the restriction enzymes according to the manufacturer's instructions. The cut solution ( $1.0\text{ }\mu\text{l}$ ) was mixed with  $12\text{ }\mu\text{l}$  of deionized formamide, and  $0.5\text{ }\mu\text{l}$  of DNA fragment length standard (GS-400HD, Applied Biosystems). The mixture was denatured at  $95^{\circ}\text{C}$  for 2 min and chilled on ice prior to electrophoresis. The fluorescently labeled terminal restriction fragments (T-RFs) were analyzed by electrophoresis on an ABI PRISM 310 Genetic Analyzer (Applied Biosystems) in GeneScan mode using the default 50 fluorescence units. Electrophoresis conditions were  $60^{\circ}\text{C}$ , 15 kv and 20 s for the injection time, with a run time of 30 min using POP-4 polymer. T-RFLP electropherograms were analyzed with GeneScan 2.1 software (Applied Biosystems).

**Numerical analysis.** Differences in T-RFLP patterns among samples were evaluated using similarity values. The similarity values were obtained by the correlation coefficient of Pearson. A dendrogram based on the similarity value's matrix was constructed using the group-mean joining methods. The horizontal scale ("rescaled distance cluster combine") used arbitrary units from 0 to 25 in order to measure the similarity between the various samples that were clustered. All methods mentioned above were conducted using SPAA version 12.0J.

## Results and Discussion

### *Diurnal changes of rumen fluid component after feeding*

Slight drops (0.6, 0.8, and 1.3) in the pH levels in rumen fluids of sheep (A, B, and C) fed the roughage diet were observed from pre-feeding values (6.6, 6.7, and 7.4) to the lowest values (6.0, 5.9, and 6.1) at 12, 16, and 6 hr after feeding, respectively (Fig. 1). Most of the pH values were well within what is considered normal (6.1-7.2) (Leedle et al. 1982) in our study, when the animals were fed the roughage diet. Generally, when animals are fed a diets containing a large proportion of readily fermentable carbohydrates, a considerable depressions in pH values will result (Mackie and




Fig. 1 Diurnal changes in pH and ammonia levels in rumen fluid from sheep fed both the roughage and the high-concentrate diet.  
sheep A (◆), sheep B (■), sheep C (▲).

Gilchrist, 1979; Mackie et al., 1978). In the present study, where the animals were fed the high-concentrate diet, remarkable decreases (2.4, 2.3, and 2.2) in the pH levels in rumen fluids of sheep (A, B, and C) were also found, that is, the pH levels changed from the pre-feeding values (7.5, 7.5, and 7.3) to the lowest values (5.1, 5.2 and 5.1) at 6, 3 and 6 hrs after feeding, respectively (Fig. 1).

Leedle et al. (1982) reported that ammonium concentration in rumen fluids of cattle fed high-forage diets increased temporarily after feeding and later decreased, and the concentration of rumen fluids fed a high-concentration diet decreased gradually after feeding. In the present study, similar diurnal variations in ammonia concentrations were observed in both diets (Fig. 1). Ammonia concentrations of rumen fluids in sheep (A, B, and C) fed the roughage diet increased from pre-feeding values of 7.6, 10.2, and 9.5 mM to maximum values of 10.8, 13.2, and 11.0 mM at 4, 2, and 2 hr after feeding, and then leveled down to minimum values of 5.8, 5.2, and 3.1 mM at 8, 14, and 8 hr after feeding, respectively. Ammonia concentrations of rumen fluids in sheep (A, B, and C) fed the high-concentration diet decreased continuously from pre-feeding values of 14.3, 10.8, and 13.6 mM to minimum values of 6.2, 1.9 and 4.4

mM at 12, 6, and 6 hr after feeding, and then leveled up in all sheep (Fig. 1).

Table 1 and 2 show total volatile fatty acid (VFA), mol% of each VFA/total VFA, and acetate/propionate (A/P) ratio in rumen fluids of sheep fed both the roughage diet and the high-concentration diets, respectively. Total VFA of rumen fluids in sheep (A, B, and C) fed the roughage diet ranged from pre-feeding values of 17.6, 27.3 and 24.5 mM, to the maximum values of 32.4, 33.4, and 33.2 mM at 12, 8, and 8 h after feeding, respectively. Total VFA of rumen fluids in animals (A, B, and C) fed the high-concentration diet ranged from pre-feeding values of 17.4, 24.3, and 18.1 mM, to the maximum values of 36.8, 46.2, and 67.5 mM at 12, 4, and 6 h after feeding, respectively. The increase of total VFA after feeding was greater in the rumen fluids of sheep fed on the high-concentration diet than in those of sheep fed the roughage diet (especially in sheep C).

In the roughage diet, acetate constituted between 48.8 and 69.7 molar % of the total acids. Propionate ranged from 20.6 to 30.2 molar % and butyrate ranged from 7.4 to 21.1 molar %. The A/P ratios calculated from these data were from 1.6 to 3.0, and became the lowest values (1.6- 2.3) at 4 hr after feeding. In the high-concentration

**Table 1** Diurnal variations in fermentation acids in ruminal fluid from animals fed on the roughage diet.

Sheep No.	Hours after feeding	Total VFA (mM)	VFA composition (mol%) <sup>a</sup>						C <sub>2</sub> /C <sub>3</sub>
			C <sub>2</sub>	C <sub>3</sub>	i-C <sub>4</sub>	n-C <sub>4</sub>	i-C <sub>5</sub>	n-C <sub>5</sub>	
Sheep A	0	17.6	57.3	25.7	2.4	8.8	4.3	1.5	2.2
	4	31.8	48.8	30.2	1.1	15.5	0.7	3.6	1.6
	8	31.9	53.4	29.4	0.6	14.5	–	2.2	1.8
	12	32.4	53.7	24.3	0.2	21.1	–	0.8	2.2
	18	24.7	64.1	26.9	0.5	7.6	–	0.9	2.4
	24	30.6	58.8	20.6	0.8	19.1	–	0.7	2.9
Sheep B	0	27.3	64.5	22.5	0.6	9.8	1.5	1.1	2.9
	4	32.2	57.6	24.6	1.6	13.3	1.0	2.0	2.3
	8	33.4	62.7	24.0	0.5	11.2	–	1.5	2.6
	12	31.7	63.7	23.6	0.6	10.8	–	1.3	2.7
	18	30.5	69.7	21.6	0.7	7.4	–	0.7	3.2
	24	27.1	67.3	22.1	1.2	7.6	0.9	0.9	3.0
Sheep C	0	24.5	65.8	24.7	0.8	7.6	–	1.1	2.7
	4	30.5	55.2	28.9	0.8	12.9	–	2.3	1.9
	8	33.2	56.2	29.0	0.9	12.3	–	1.5	1.9
	12	28.5	58.4	28.6	0.5	10.9	–	1.6	2.0
	18	26.5	57.7	27.6	2.0	9.9	1.3	1.5	2.1
	24	18.6	57.9	27.5	3.6	9.8	–	1.2	2.1

<sup>a</sup> C<sub>2</sub>, acetic acid; C<sub>3</sub>, propionic acid; i-C<sub>4</sub>, iso-butyric acid; n-C<sub>4</sub>, n-butyric acid; i-C<sub>5</sub>, iso-valeric acid; n-C<sub>5</sub>, n-valeric acid.

**Table 2** Diurnal variations in fermentation acids in ruminal fluid from animals fed on the high-concentrate diet.

Sheep No.	Hours after feeding	Total VFA (mM)	VFA composition (mol%) <sup>a</sup>							C <sub>2</sub> /C <sub>3</sub>
			C <sub>2</sub>	C <sub>3</sub>	i-C <sub>4</sub>	n-C <sub>4</sub>	i-C <sub>5</sub>	n-C <sub>5</sub>	n-C <sub>6</sub>	
Sheep A	0	17.4	51.0	18.3	3.3	17.2	7.8	2.5	–	2.8
	2	30.6	37.8	31.1	2.3	17.2	1.7	9.0	0.9	1.2
	4	33.8	49.4	20.9	0.8	23.2	2.1	3.6	–	2.4
	6	28.9	48.5	17.7	0.8	25.7	3.1	3.9	0.3	2.7
	12	36.8	50.9	19.3	0.5	23.3	2.5	3.5	–	2.6
	18	23.9	49.6	19.3	1.8	21.2	5.0	3.1	–	2.6
	24	17.6	54.0	20.2	2.5	17.0	4.3	2.1	–	2.7
Sheep B	0	24.3	43.2	31.0	3.4	9.4	4.8	7.6	0.6	1.4
	2	32.0	47.6	25.0	2.4	19.5	2.5	2.9	–	1.9
	4	46.2	35.0	25.1	1.2	28.3	0.5	8.6	1.4	1.4
	6	40.2	40.5	23.4	0.4	23.9	0.5	9.0	2.4	1.7
	12	33.3	43.4	27.2	0.9	22.1	1.4	0.9	4.1	1.6
	18	31.7	32.5	26.5	1.8	9.0	3.3	7.2	19.8	1.2
	24	18.9	40.4	39.4	3.1	5.5	5.3	6.4	–	1.0
Sheep C	0	18.1	44.7	34.9	3.0	6.6	5.4	5.1	0.4	1.3
	2	44.9	32.2	24.5	1.7	28.2	1.2	10.2	2.1	1.3
	4	59.9	31.2	18.5	1.2	32.4	0.4	13.0	3.4	1.7
	6	67.5	29.8	14.0	0.3	33.9	0.4	15.5	6.1	2.1
	12	60.0	35.3	17.0	0.6	22.7	0.6	14.0	9.8	2.1
	18	31.5	36.2	26.9	2.5	13.8	4.4	12.3	3.8	1.3
	24	21.7	43.1	35.8	0.8	9.1	5.0	6.0	0.3	1.2

<sup>a</sup>Some as Table 1.

diet, acetate constituted between 29.8 and 54.0 molar % of the total acids. Propionate and butyrate constituted 14.0–39.4 and 5.5–33.9 molar %, respectively. The A/P ratios were from 1.0 to 2.8. The extent of changes in total VFA concentration, and molar % of acetate, pro-

pionate, iso-valerate, and valerate, each, in the rumen fluids of sheep fed the high-concentrate diet, were greater than in those fed the roughage diet in all sheep. Iso-caproate was detected only when the high-concentrate diet was given. When pH in the rumen becomes lower than 5.5 when

feeding on a high-grain diet, lactic acid accumulates in the rumen (Nocek, 1997). Although the pH of rumen fluids of sheep fed the high-concentrate diet became less than 5.5 in some cases, lactate (0.5 mM) was detected only in the rumen fluid of sheep A at 2 hr after feeding in the present study (data not shown). In conclusion, serious diurnal variations of pH,  $\text{NH}_4$  concentration and VFA concentration and VFA % were found in rumen fluids of sheep fed the high-concentrate diet. More so than in those of the sheep fed the roughage diet after feeding.

#### *Selection of restriction enzyme*

T-RFLP analysis was applied to the bovine rumen sample obtained from the slaughterhouse mentioned previously, and T-RFLP fingerprinting was compared with three tetrameric restriction enzymes. *Hha*I generated 13 terminal restriction fragments (T-RFs), *Msp*I generated 25 T-RFs, and *Afa*I generated 6 T-RFs within a size of less than 400 bp (Fig. 2). According to computer simulations of the sequences in the Ribosomal Database Project small-subunit database (Maidak et al., 1996), *Msp*I was one of the restriction enzymes that produced the largest number of unique T-RFs (Liu et al., 1997). As digestion with *Msp*I

resulted in the largest number of T-RFs in our experiment, and it was suggested that the enzyme could provide considerable elucidation in demonstrating the differences in the bacterial communities between the samples, only *Msp*I was used in the next analyses.

#### *Diurnal changes on T-RFLP patterns after feeding*

Before feeding, similar R-RFLP patterns between the three sheep were recognized in the roughage diet (RA-0, RB-0, and RC-0), however some animal-to-animal variations in the patterns was found concerning the high-concentration diet (CA-0, CB-0, and CC-0). The representative T-RFLP profiles (RB-0 and CB-0 in sheep B) before being fed the roughage and the high-concentration diets are presented in Fig. 3.

The diurnal changes of T-RFLP profiles were examined for both the roughage and the concentrate diets. In the roughage diet, a total of 69 fragments (sheep A, 59; sheep B, 52; sheep C, 38 fragments, each) were detected, and in the high-concentrate diet a total of 64 fragments (sheep A, 26; sheep B, 38; sheep C, 36 fragments) were detected (Table 3 and 4). In total, 101 fragments were included in the analyses. In the experiment





Fig. 2 T-RFLP analyses corresponding to the digestion with the restriction enzymes of *Hha*I, *Msp*I, and *Afa*I.



**Fig. 3** Comparison of T-RFLP patterns of 16S rDNA from rumen bacterial fraction fed the roughage diet (upper) and the concentrate diets (below) in sheep B.

concerning the roughage diet, the number of fragments detected was between 28 and 35 in sheep A, 24 and 34 in sheep B, and between 19 and 31 in sheep C, respectively. In the experiment employing the high-concentrate diet, the number of fragments detected was between 7 and 16 in sheep A, 10 and 28 in sheep B, and between 14 and 23 in sheep C, respectively. Thirty-two T-RFs (89, 93, 95, 121, 125, 145-147, 158, 186, 210, 213, 219, 221, 238, 273, 279-280, 282, 286-288, 291-292, 294-297, 310-311, and 332-333 bp) were universally distributed, i.e., they were found in all samples, while others varied in distribution and were related to specific diets. Thirty-seven fragments (82, 86-87, 91, 96, 123-124, 131, 134-136, 143, 156, 164, 166, 169, 209, 211-212, 214, 218, 224-225, 233-234, 266, 272, 276-277, 281, 283-285, 301, 305-306 and 313 bp) were found only in animals receiving the roughage diet, and 32 fragments (32-39, 44-45, 47, 90, 92, 116, 128, 144, 155, 160, 163, 190-191, 201, 220, 240, 248, 252, 267, 270, 278, 289, 303 and 309 bp) were only found in animals which received the high-concentrate diet. Bacterial communities with different species compositions produce different and characteristic T-RFLP profiles. Diet type greatly influenced the rumen bacterial community, as is expressed by the T-RFLP profiles in the present

study.

A dendrogram (Fig. 4) was constructed by the group-mean joining method on the basis of the similarity matrix data (data not shown). Separated clusters were formed between the roughage diet group and the high-concentrate diet group. Rumen samples from sheep fed the roughage diet formed a close cluster having distances less than 9 units. The rumen samples from sheep fed the high-concentrate diet, sheep A and B, belonged to the same group having distances of less than 22 units. The samples from sheep C fed the concentrate diet, formed another group having distances of less than 25 units against the samples from these sheep. Judging from such differences of distance in the dendrogram, it was presumed that the bacterial communities in the rumen were rather stable among the animals being fed the roughage diet, but the communities were considerably destabilized in the animals being fed the high-concentration diet.

About twelve bands generated from rumen bacterial communities of roughage or corn diets by DGGE analysis (Kocherginskaya et al., 2001). Since a total of 69 T-RFs (roughage diet) or 64 (concentrate diet) were detected in our study, T-RFLP analysis seemed to be more useful than



**Table 3** Terminal restriction fragments included in the analyses of rumen bacterial fraction from animals fed on the roughage diet.

RFs (bp)	Sheep A					Sheep B					Sheep C							
	RA-0 <sup>1)</sup>	RA-4	RA-8	RA-12	RA-18	RA-24	RB-0	RB-4	RB-8	RB-12	RB-18	RB-24	RC-0	RC-4	RC-8	RC-12	RC-18	RC-24
82	0.6	0.7	0.7	0.6	0.9	0.7	-	-	-	0.7	-	-	-	-	-	-	-	-
86	0.6	-	-	-	0.5	1.6	-	0.7	1.7	2.6	1.5	1.6	2.5	2.2	2.2	2.5	2.0	2.0
87	1.1	1.4	0.8	1.9	2.4	-	1.4	1.1	1.7	9.8	13.9	12.4	9.4	13.0	13.6	17.2	15.9	14.3
89	11.4	11.9	11.3	17.4	14.9	12.0	13.5	14.6	9.8	14.0	16.6	12.4	9.4	13.0	13.6	17.2	15.9	14.3
91	1.7	1.5	1.0	1.2	1.3	1.4	1.2	1.7	1.8	1.6	1.9	1.9	1.5	1.2	1.2	1.5	2.1	1.7
93	14.4	20.6	7.9	26.2	24.8	24.3	20.8	20.5	23.3	16.6	27.0	23.8	18.6	30.7	27.0	28.8	16.6	28.0
95	1.2	0.6	-	1.9	2.6	2.9	1.4	1.7	3.1	1.3	-	3.5	2.6	2.6	2.8	2.8	-	2.9
96	0.6	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
121	2.0	1.7	0.7	0.9	0.6	1.2	0.7	1.1	1.4	0.6	-	-	0.8	-	1.4	-	-	1.5
123	-	-	-	-	-	-	0.8	0.9	-	-	1.3	2.2	1.0	-	-	-	-	-
124	2.4	1.2	2.2	3.6	3.7	2.0	3.5	3.7	1.7	3.8	3.4	2.6	2.2	2.2	2.5	2.7	-	1.9
125	5.4	3.1	1.8	1.0	1.1	1.1	1	2.2	2.1	0.9	-	-	1.0	-	1.1	-	2.4	2.3
131	-	-	-	0.7	0.6	-	-	-	-	-	-	-	-	-	-	-	-	-
134	-	-	-	0.7	0.8	0.7	1.2	-	-	-	1.2	1.6	1.0	1.0	0.8	-	-	-
135	-	-	-	-	1.1	0.9	-	1.0	1.0	0.7	-	-	-	-	0.8	-	-	-
136	-	-	-	0.7	-	-	0.5	0.7	-	-	-	-	-	-	0.8	-	-	1.5
143	1.2	0.6	0.6	0.9	-	-	-	0.7	-	1.0	-	-	-	-	-	-	-	-
145	-	-	0.6	0.6	0.9	-	0.6	0.6	0.8	0.8	-	-	-	-	-	-	-	-
146	1.5	1.8	1.1	1.1	2.1	2.0	2.8	2.3	2.8	2.0	2.8	3.2	3.0	1.4	1.0	0.9	-	1.5
147	-	-	-	-	2.5	2.4	-	-	-	-	-	-	-	3.6	4.3	2.9	-	1.2
156	0.7	0.8	0.6	-	-	0.6	-	0.7	-	-	-	-	-	-	-	-	-	-
158	1.2	0.9	0.6	-	-	-	0.5	0.7	-	-	-	-	-	-	-	-	-	-
164	0.5	0.6	-	-	-	-	0.6	0.7	-	0.6	-	-	-	-	-	-	-	-
166	0.8	-	-	-	-	-	0.6	0.7	-	-	-	-	-	-	-	-	-	-
186	-	-	-	-	0.8	1.1	0.8	0.9	0.8	1.3	1.4	1.4	0.9	-	0.7	-	-	-
189	0.9	-	-	1.2	-	1.7	-	-	-	-	-	-	-	-	1.1	0.8	-	-
209	-	-	-	-	-	0.9	-	-	-	-	-	-	-	-	-	-	-	-
210	-	-	-	-	-	0.9	-	-	-	-	-	-	-	-	-	-	-	-
211	-	-	-	-	0.5	-	-	-	-	-	-	-	-	-	-	-	-	-
212	-	-	-	-	-	0.9	-	-	-	-	-	-	-	-	-	-	-	-
213	0.7	-	-	1.8	2.4	0.7	-	-	0.8	-	-	0.9	-	1.3	1.0	1.1	-	-
214	0.9	1.6	1.5	1.2	-	-	1.2	1.8	0.9	1.7	1.9	1.3	-	1.3	1.5	1.7	-	1.3
218	-	-	0.8	-	-	0.7	1.8	2.5	-	2.3	-	1.0	-	-	-	-	-	-
219	1.5	1.5	0.9	0.8	0.6	1.5	-	-	1.4	-	1.3	-	1.1	1.0	1.5	1.1	-	-
221	3.9	3.1	4.4	2.0	2.3	0.6	4.1	2.8	5.0	3.2	4.9	4.1	4.3	1.8	2.0	2.1	4.7	3.8
224	-	2.2	-	-	-	-	-	-	1.4	-	-	-	-	-	-	-	-	-
225	1.5	1.5	1.7	-	-	-	0.8	1.2	1.6	-	-	-	0.8	-	-	-	-	-
233	8.0	4.9	18.5	3.6	12.3	-	12.0	11.2	12.1	5.6	-	5.7	6.8	-	4.2	3.6	2.2	1.6
234	-	5.6	-	6.4	-	-	6.4	-	-	6.4	13.1	7.8	7.7	13.2	7.6	6.3	11.3	7.0
238	-	-	0.8	0.6	1.1	0.9	-	-	-	-	-	1.1	0.6	-	-	6.7	11.4	6.7
266	-	-	-	0.8	-	-	-	-	-	-	1.2	-	-	2.4	2.6	2.0	-	-
272	-	-	-	-	-	-	-	-	-	-	0.8	-	-	-	-	-	-	-
273	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
276	-	-	-	-	0.6	-	-	-	-	-	-	-	-	-	-	-	-	-
277	-	-	-	-	-	-	-	-	0.8	-	1.1	-	0.8	-	-	-	-	-
279	-	-	-	-	2.1	-	-	-	-	-	-	-	-	-	-	-	-	-
280	-	-	-	-	4.2	1.2	3.7	1.1	-	2.2	1.5	5.0	6.8	-	-	-	2.2	2.7
281	0.6	0.9	1.7	2.8	-	-	-	-	-	1.5	-	-	-	2.2	1.3	2.2	2.0	-
282	-	0.5	-	-	-	1.2	1.8	0.8	1.3	1.5	-	1.1	1.7	-	-	-	-	-
283	-	-	0.7	1.5	-	2.9	1.1	0.6	-	1.4	0.9	1.5	1.9	1.4	1.2	1.1	1.9	-
284	-	-	-	-	-	-	1.1	0.6	-	-	-	1.0	1.9	1.0	-	-	-	-
285	-	1.2	-	-	-	-	0.6	-	-	2.8	-	-	-	-	-	1.5	4.4	-
286	2.3	1.3	4.4	1.1	-	-	-	-	2.7	-	-	-	-	-	-	-	-	1.7
287	-	-	-	-	-	1.3	-	-	-	-	-	-	-	-	-	-	-	-
288	1.2	1.6	1.3	-	-	-	0.8	0.7	1.1	1.3	-	-	0.9	-	-	-	1.8	-
291	0.7	0.9	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
292	1.9	2.1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
294	-	-	-	-	-	-	3.5	-	4.1	4.7	-	2.2	3.6	-	2.0	1.9	3.2	-
295	-	-	-	2.5	-	-	-	-	-	-	6.0	-	-	-	-	-	-	-
296	6.4	7.4	7.0	3.4	4.6	-	2.7	9.3	3.9	5.1	-	2.5	3.0	-	2.8	5.9	4.9	5.9
297	6.6	4.6	6.0	-	-	7.9	3.6	3.3	5.9	3.1	3.7	2.6	3.4	5.6	3.1	-	5.7	4.1
301	-	-	-	-	-	1.0	-	-	-	-	-	-	-	-	-	-	-	-
305	12.1	8.0	15.6	6.3	3.5	1.1	-	-	2.3	-	-	-	-	-	5.7	6.1	-	-
306	-	-	-	-	-	-	2.8	3.4	-	4.6	4.5	3.5	3.5	-	-	-	3.5	2.4
310	2.4	2.2	3.5	1.7	2.1	-	3.0	2.0	2.7	1.9	1.9	1.9	3.1	2.1	1.7	1.7	2.0	-
311	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
313	1.0	0.8	1.7	1.8	3.4	3.8	3.4	1.7	2.1	1.0	2.1	1.5	3.0	2.2	1.4	1.2	-	1.8
332	-	-	-	0.7	0.8	-	-	0.6	-	-	0.9	0.9	-	-	-	-	-	-
333	-	-	-	-	-	0.7	-	-	-	-	-	-	-	-	-	-	-	-

Values are presented as percentage of total areas.

<sup>1)</sup> A, B, and C, sheep No.; The final number, hours after feeding.

**Table 4** Terminal restriction fragments included in the analyses of rumen bacterial fraction from animals fed on the concentrate diet.

RFs (bp)	Sheep A					Sheep B					Sheep C				
	CA-0''	CA-6	CA-12	CA-18	CA-24	CB-0	CB-6	CB-12	CB-18	CB-24	CC-0	CC-6	CC-12	CC-18	CC-24
32	-	-	-	2.4	1.7	4.6	1.5	1.0	-	-	0.6	0.8	-	1.6	-
33	-	-	4.7	5.4	3.7	4.9	1.3	2.3	4.4	6.6	1.4	2.3	1.7	2.1	1.6
34	2.6	2.2	6.9	6.2	4.4	4.1	-	3.1	6.6	-	1.0	2.9	-	-	-
35	3.5	-	-	-	-	-	1.5	-	-	4.7	-	-	-	-	1.5
36	2.9	2.8	7.0	5.1	5.1	2.3	2.5	2.9	6.5	-	1.6	2.2	-	1.0	3.0
37	2.2	2.6	6.7	3.9	3.9	1.6	1.1	-	7.2	-	1.1	2.8	-	-	2.7
38	-	-	-	-	-	-	-	-	2.5	-	-	-	-	-	-
39	-	-	-	-	-	0.8	-	0.7	-	-	-	-	-	-	-
44	-	-	-	-	-	1.3	-	-	-	-	-	-	-	-	-
45	-	-	-	-	-	2.4	-	-	2.6	-	-	-	-	-	-
47	-	-	-	-	1.3	-	-	-	-	-	-	-	-	-	-
89	-	-	-	-	-	1.8	-	-	-	2.3	-	-	-	-	1.6
90	-	-	-	-	-	-	-	-	-	3.3	-	-	-	-	-
92	-	-	-	-	-	2.1	-	-	-	-	-	-	-	-	-
93	-	-	-	-	2.2	2.6	0.8	-	-	2.7	0.7	-	7.3	2.7	-
95	-	2.4	-	-	-	16.7	9.9	-	-	2.2	-	-	-	-	-
116	-	-	-	-	-	-	-	-	-	1.3	-	-	-	-	-
121	3.6	6.6	-	-	-	4.9	3.4	0.9	-	6.1	1.9	-	1.8	-	1.4
125	5.9	10.4	-	-	1.3	2.8	3.5	0.9	-	3.2	-	-	-	-	-
128	-	-	-	-	-	-	0.7	-	-	-	-	-	-	-	-
144	-	-	-	-	-	-	-	58.5	-	3.2	-	-	-	-	-
145	-	3.8	-	-	-	7.9	24.6	-	7.2	5.0	-	-	1.1	-	-
146	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
147	-	-	-	-	-	1.0	-	-	-	-	0.6	-	-	-	-
155	-	-	-	-	-	0.8	0.9	-	-	-	-	-	-	-	-
158	-	1.7	-	-	-	0.5	-	-	-	-	-	-	-	-	-
160	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
163	-	-	-	3.9	9.4	-	-	-	-	-	0.6	-	-	-	-
186	-	-	18.7	17.8	-	-	1.0	0.7	3.5	3.5	-	2.3	-	-	15.7
190	-	-	-	-	-	-	-	-	-	-	27.0	14.4	25.6	31.8	-
191	-	-	-	-	-	1.3	1.0	1.1	-	2.6	-	1.1	-	-	-
201	-	-	-	-	-	-	-	-	-	-	1.3	-	-	-	-
210	-	-	-	-	-	-	-	-	-	-	0.9	-	1.6	-	-
213	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
219	-	-	-	-	-	-	6.0	4.1	-	-	-	-	-	-	1.3
220	-	-	-	-	-	-	8.2	-	-	-	-	-	2.1	3.6	5.5
221	-	-	-	-	-	1.6	-	-	-	10.6	-	-	-	2.0	2.0
238	3.2	-	-	-	-	-	-	-	-	2.6	9.4	4.3	5.6	6.7	5.3
240	-	-	-	-	-	-	-	-	-	-	1.0	-	-	-	-
248	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
252	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
267	-	5.2	-	-	-	-	-	-	-	-	-	-	-	-	-
270	15.5	4.7	47.9	47.2	8.5	-	2.8	-	-	-	-	-	-	-	-
273	-	-	-	-	-	0.9	-	10.2	48.6	6.3	-	-	-	-	-
278	-	-	-	-	-	-	-	-	-	3.2	-	-	-	-	-
279	-	-	-	2.0	-	-	-	-	-	-	8.3	5.0	2.4	8.1	3.5
280	-	-	-	-	1.6	1.4	-	-	-	-	-	-	-	-	-
282	2.8	-	-	-	-	-	-	-	-	-	-	3.0	-	-	-
286	-	-	-	-	-	-	-	-	-	-	3.6	28.9	5.4	-	-
287	-	-	-	-	-	-	-	-	-	-	5.5	-	19.4	18.0	17.1
288	-	2.9	-	-	-	4.0	3.3	-	-	10.2	-	-	-	-	-
289	-	5.2	-	-	-	-	-	-	-	-	2.1	1.5	3.3	-	1.0
291	-	-	-	-	-	1.6	0.7	-	-	-	-	-	-	-	-
292	4.9	4.6	-	-	2.4	1.4	1.6	-	-	-	-	-	-	-	-
294	-	-	-	-	-	-	1.1	-	-	-	-	-	-	1.3	-
295	-	-	-	-	-	-	-	-	-	-	1.2	-	-	-	-
296	20.7	9.8	8.0	6.0	52.8	8.4	19.7	13.5	10.8	20.3	1.6	6.2	4.9	1.2	3.4
297	-	-	-	-	-	-	-	-	-	-	16.2	10.4	8.7	1.9	-
303	32.3	31.4	-	-	1.9	9.1	3.0	-	-	-	-	-	2.8	10.4	11.6
309	-	-	-	-	-	-	-	-	-	-	-	10.8	-	-	1.3
310	-	3.7	-	-	-	-	-	-	-	-	-	-	-	-	-
311	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1.1
332	-	-	-	-	-	-	-	-	-	-	-	-	2.1	-	-
333	-	-	-	-	-	-	-	-	-	-	3.4	1.3	4.3	9.4	19.4

Values are presented as percentage of total areas.

<sup>1)</sup> Same as table 3.




Fig. 4 A dendrogram made from T-RFLP analyses of the roughage and the concentrate diets. The symbols and the number are the same as in table 3 and table 4.

DGGE analysis in comparing rumen bacterial communities.

T-RF lengths can be predicted from known 16S rRNA gene sequences, and are available as supplementary data in JMM Online (<http://jmm.sgmjournals.org>). But it is difficult to relate each of the T-RFs to a specific bacterial group, because in many cases a single T-RF can represent several genera and the discrepancies between predicted and observed T-RF lengths have been reported previously (Clement et al., 1998; Sakamoto et al., 2003). To clarify the bacterial origin of each T-RF, we performed T-RFLP analyses for stoke cultures in our laboratory (reported elsewhere). 93 and 296 bp presumed to be the T-RFs derived from *Prevotella* sp., and *Selenomonas ruminantium*, respectively, from the analyses. The authors believe that T-RFLP analysis of pure culture is required for accurate identification of each T-RF. A more accurate

community representative may be achieved by combining T-RFLP data derived from the digestion of different enzymes (Clement et al., 1998) or the primer-enzyme combinations for T-RFLP analysis (Nagashima et al., 2003). Many T-RFs remained in area more than 400 bp (Fig. 2). Extension to a range more than 400 bp in the analysis is needed to grasp the difference or the change of rumen bacterial composition in detail. In conclusion, T-RFLP analysis is useful for demonstrating changes in the bacterial community structure in sheep fed different experimental diets. It may be exploited for rapid detection of other rumen microbial changes such as those imposed by disease, addition of antibiotics, or lactic acidosis development.

## References

- Attwood, G. T., Lockington, R. A., Xue, G. P., and Brooker, J. D. (1988) Use of a unique gene sequence as a probe to enumerate a strain of *Bacteroides rumenicola* introduced in to the rumen. *Appl. Environ. Microbiol.*, **54**, 534-539.
- Avaniss-Aghajani, E., Jones, K., Chapmen, D., and Brunk, C. (1994) A molecular technique for identification of bacteria using small subunit ribosomal RNA sequences. *BioTechniques*, **17**, 144-149.
- Blackwood, C. B., Marsh T., Kim S.-H., and Paul E. A. (2003) Terminal restriction fragment length polymorphism data analysis for quantitative comparison of microbial communities. *Appl. Environ. Microbiol.*, **69**, 926-932.
- Brosius, J., Palmer, J. L., Kennedy, J. P., and Noller, H. F. (1978) Complete nucleotide sequence of 16S ribosomal RNA gene from *Escherichia coli*. *Proc. Natl. Acad. Sci. USA*, **75**, 4801-4805.
- Chaney, A. L. and Marbach, E. P. (1962) Modified reagents for determination of urea and ammonia. *Clin. Chem.*, **8**, 130-132.
- Clement, B. G., Kehl, E., Debord, K. L., and Kitts, C. L. (1998) Terminal restriction fragment patterns (TRFPs), a rapid, PCR-based method for the comparison of complex bacterial communities. *J. Microbiol. Method*, **31**, 135-142.
- Dehority, B. A. and Orpin, C. G. (1977) Develop-

- ment of, and natural fluctuations in, rumen microbial populations. In *The Rumen Microbial Ecosystem*, ed. by Hobson, P. N. and Stewart, C. S., Chapman & Hall, London, pp. 196-245.
- Dunbar, J., Ticknor L. O., and Kuske C. R. (2000) Assessment of microbial diversity in four south-western united states soils by 16S rRNA gene terminal restriction fragment analysis. *Appl. Environ. Microbiol.*, **66**, 2943-2950.
- Hayashi, H., Sakamoto M., and Benno Y. (2002) Fecal microbial diversity in a strict vegetarian as determined by molecular analysis and cultivation. *Microbiol. Immunol.*, **46**, 819-831.
- Hiraishi, A., Iwasaki M., and Shinjo H. (2000) Terminal restriction pattern analysis of 16S rRNA genes for the characterization of bacterial communities of activated sludge. *J. Biosci. Bioeng.*, **90**, 148-156.
- Kitts, C. L. (2001) Terminal restriction fragment patterns: a tool for comparing microbial communities and assessing community dynamics. *Curr. Issues. Intest. Microbiol.*, **2**, 17-25.
- Kocherginskaya, S. A., Aminov R. I., and White B. A. (2001) Analysis of the rumen bacterial diversity under two different diet conditions using denaturing gradient gel electrophoresis, random sequencing, and statistical ecology approaches. *Anaerobe*, **7**, 119-134.
- Koike, S., Pan, J., Kobayashi, Y., and Tanaka, K. (2003) Kinetics of *in sacco* fiber-attachment of representative ruminal cellulolytic bacteria monitored by competitive PCR. *J. Dairy Sci.*, **86**, 1429-1435.
- Lane, D. J. (1991) 16S/23S rRNA sequencing. In *Nucleic Acid Techniques in Bacterial Systematics*, ed. by Stackebrandt, E. and Goodfellow, M., Wiley, Chichester, pp. 115-175.
- Leedle, J. A. Z., Bryant, M. P., and Hespell, R. B. (1982) Diurnal variations in bacterial numbers and fluid parameters in ruminal contents of animals fed low- or high-forage diets. *Appl. Environ. Microbiol.*, **44**, 402-412.
- Leser, T. D., Lindecrone, R. H., Jensen, T. K., Jensen, B. B., and Møller K. (2000) Changes in bacterial community structure in the colon of pigs fed different experimental diets and after infection with *Brachyspira hyodysenteriae*. *Appl. Environ. Microbiol.*, **66**, 3290-3296.
- Liu, W. -T., Marsh T. L., Chen H., and Forney L. J. (1997) Characterization of microbial diversity by determining terminal restriction fragment length polymorphisms of genes encoding 16S rRNA. *Appl. Environ. Microbiol.*, **63**, 4516-4522.
- Mackie, R. I. and Gilchrist, M. C. (1979) Changes in lactate-producing and lactate-utilizing bacteria in relation to pH in the rumen of sheep during stepwise adaptation to a high-concentrate diet. *Appl. Environ. Microbiol.*, **38**, 422-430.
- Mackie, R. I., Gilchrist, F. M. C., Roberts, A. M., Hannah, P. E., and Schwartz, H. M. (1978) Microbiological and chemical changes in the rumen during stepwise adaptation of sheep to high concentrate diets. *J. Agric. Sci.*, **90**, 241-254.
- Maidak, B. L., Olsen, G. J., Larsen, N., Overbeek, R., McCaughey, M. J., and Woese, C. R. (1996) The Ribosomal Database Project (RDP). *Nucleic Acids Res.*, **24**, 82-85.
- Minato, H., Miyagawa, E., and Suto T. (1990) Technique for analysis of rumen microbial ecosystems. In *The Rumen Ecosystem*, ed. Hoshino, S., Onodera, R., Minato, H., and Itabashi, H., Japan Scientific Societies Press, Tokyo, pp. 3-12.
- Nagashima, K., Hisada. T., Sato, M., and Mochizuk, J. (2003) Application of new primer-enzyme combinations to terminal restriction fragment length polymorphism profiling of bacterial populations in human feces. *Appl. Environ. Microbiol.*, **69**, 1251-1262.
- Nocek, J. E. (1997) Bovine acidosis: implication on laminitis. *J. Dairy Sci.*, **80**, 1005-1028.
- Sakamoto, M., Takeuchi, Y., Umeda, M., Ishikawa, I., and Benno, Y. (2003) Application of terminal RFLP analysis to characterize oral bacterial flora in saliva of healthy subjects and patients with periodontitis. *J. Med. Microbiol.*, **52**, 79-89.
- Scala, D. J. and Kerkhof, L. J. (2000) Horizontal heterogeneity of denitrifying bacterial communities in marine sediments by terminal restriction fragment length polymorphism analysis.

- Appl. Environ. Microbiol.*, **66**, 1980-1986.
- Stahl, D. A., Flesher, B., Mansfield, H. R. and Montgomery, L. (1988) Use of phylogenetically based hybridization probes for studies of ruminal microbial ecology. *Appl. Environ. Microbiol.*, **54**, 1079-1084.
- Suto, T. (1970) Ecological studies on the rumen bacteria and anaerobes. *Nat. Inst. Anim. Hlth Quart.*, **10**, Suppl., 109-130.
- Tajima, K., Aminov, R. I., Nagamine, T., Matsui, H., Nakamura, M., and Benno, Y. (2001) Diet-dependent shifts in the bacterial population of the rumen revealed with real-time PCR. *Appl. Environ. Microbiol.*, **67**, 2766-2774.
- Tajima, K., Aminov, R. I., Nagamine, T., Ogata, K., Nakamura, M., Matsui, H., and Benno, Y. (1999) Rumen bacterial diversity as determined by sequence analysis of 16S rDNA libraries. *FEMS Microbiol. Ecol.*, **29**, 159-169.
- Ueki, A., Miyagawa E., Minato H., Azuma R., and Suto T. (1978) Enumeration and isolation of anaerobic bacteria in sewage digester fluids. *J. Gen. Appl. Microbiol.*, **24**, 317-332.
- Whitford, M. F., Forster, R. J., Beard, C. E., Gong, J., and Teather, R. M. (1998) Phylogenetic analysis of rumen bacteria by comparative sequence analysis of cloned 16S rRNA genes. *Anaerobe*, **4**, 153-163.
- Yanagita, K., Kamagata, Y., Kawaharasaki, M., Suzuki, T., Nakamura, Y., and Minato, H. (2000) Phylogenetic analysis of methanogens in sheep rumen ecosystem and detection of *Methanomicrobium mobile* by fluorescence in situ hybridization. *Biosci. Biotechnol. Biochem.*, **64**, 1737-1742.

## 要 旨

綿羊に粗飼料のみ、あるいは濃厚飼料主体の飼料を給与した場合について、16S リボソーム RNA 遺伝子をターゲットにした末端制限酵素断片長多型 (T-RFLP) 分析によりルーメン細菌群集構造を比較した。粗飼料あるいは濃厚飼料主体の飼料を給与して10日後の飼料給与前のルーメン液についてT-RFLP分析を行ったところ、互いに異なったピークパターンを示し、飼料によるルーメン細菌群集構造の違いが観察された。ルーメン液のpH、アンモニア濃度、揮発性脂肪酸(VFA)濃度、各VFA%の日周変動は濃厚飼料主体の飼料を給与した方が、粗飼料のみを給与した場合より大きかった。飼料の違いによるルーメン細菌群集構造の日周変動について、T-RFLP分析により解析を行った。粗飼料のみを給与した場合には、飼料給与後に、ピークパターンはあまり変動しなかった。しかし、濃厚飼料主体の飼料を給与した場合には、多くのピークが変動し細菌群集構造の不安定性が示唆された。これらの結果は、T-RFLP法によりルーメン細菌群集構造の変動を迅速、簡便に分析できることを示している。

## Summary

The structure of the bacterial community in the rumen of sheep fed a roughage diet or high-concentrate diet was compared using terminal restriction fragment length polymorphism (T-RFLP) analysis targeting the 16S ribosomal DNA. After being fed the roughage or high-concentrate diet for 10 days, reciprocal differences in the bacterial community structure in sheep rumen were detected in the form of different profiles of terminal restriction fragments (T-RFs). More serious diurnal variations in pH, NH<sub>4</sub> concentration and volatile fatty acid (VFA) concentration and VFA % were found in rumen fluids of sheep fed the high-concentrate diet than in those fed the roughage diet. Diurnal variations in the bacterial community based on differences of diet, were also analyzed by T-RFLP analysis. In the roughage diet, the profiles did not change considerably after feeding. After being fed the high-concentrate diet, many of the T-RFs fluctuated, suggesting destabilization of the microbial community. These results indicate that changes in the rumen bacterial community can be analyzed promptly and efficiently using the T-RFLP method.